

CONTAINS LOW MOLECULAR WEIGHT ORGANIC ACID AND HUMECTANT

NOT FOR USE IN ORGANIC CROP AND ORGANIC FOOD PRODUCTION.

California:

CONTAINS NONPLANT FOOD INGREDIENT(S)

17%.....Lactic acid

2%......D-Glucopyranose, oligomeric, decyl octyl glycoside

81%.....Aqueous solution/Total inert ingredients (as non-plant food)

All other states:

CONTAINS NONPLANT FOOD INGREDIENT(S)

Active Ingredients, Soil/Plant Amending Ingredients:

17%.....Lactic acid

2%......D-Glucopyranose, oligomeric, decyl octyl glycoside

81%.....Agueous solution/Total inert ingredients (as non-plant food)

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

SHAKE WELL BEFORE USE KEEP OUT OF REACH OF CHILDREN DANGER





Flammable liquid and vapor. Causes severe skin burns and eve damage.

Keep away from heat/sparks/open flames/hot surfaces. No smoking. Keep container tightly closed. Ground/bond container and receiving equipment. Use explosion-proof electrical/ventilating/lightning equipment. Use only non-sparking tools. Take precautionary measures against static discharge. Do not breathe dusts or mists. Wash thoroughly after handling. Wear protective gloves and protective clothing. Wear eye protection or face protection.

FIRST AID

Call a POISON CENTER or doctor/physician if you feel unwell.

For Chemical Emergency (spill, leak, fire or accident), call CHEMTREC (800) 424-9300.

IF ON SKIN (OR HAIR): Take off immediately all contaminated clothing. Rinse skin with water/shower. Wash contaminated clothing before reuse.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a poison center/doctor if you feel unwell.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

NET CONTENTS 2.5 US GAL (9.46 L)

GUARANTEED AND MANUFACTURED BY
Oro Agri, Inc.
2788 S. Maple Ave., Fresno, CA 93725
Oro Agri, Inc. All Rights Reserved
Copyright © May 2024 v20240528



FL License #: F002559

DENSITY 8.76 lbs/gal @ 68°F

NET WEIGHT 21.90 lbs (48.28 kg)

GENERAL INFORMATION

GALACTEK™ is designed to aid in uniform germination, root development, and overcoming post-planting stress and improve nutrient availability and retention of moisture content in soil.

DIRECTIONS FOR USE

Prior to use, read all the directions on this label and on the labels of the manufacturers' products being used with GALACTEK. The manufacturers' labels for specific usage and restrictions have precedence over directions for use referred to on the GALACTEK label. Pre-rinse all tanks, spray lines or containers thoroughly before adding GAI ACTEK

APPLICATION PRECAUTIONS & COMPATIBIL TY

Do not tank mix with products favoring a neutral to high pH, e.g., copper. Do not add to fertigation water in hydroponic cropping systems. Do not mix GALACTEK with undiluted tank mix partners until you have "jar tested" in a separate container using the proportionate amount of GALACTEK to make sure the products are compatible. Agitate and let stand for 10 minutes. Settling or separation that does not disperse indicates an incompatible mixture.

GALACTEK is compatible with ROVENSA NEXT adjuvants such as WETCIT®, VINTRE®, etc. Contact a ROVENSA NEXT sales representative when mixing with plant growth regulators in the same tank.

MIXING DIRECTIONS

Fill tank with the desired amount of water and begin agitation. Add the desired amount of pesticides, herbicides, and nutrients, if applicable. Add the desired amount of GALACTEK. Add the desired amount of adjuvants, if recommended.

USF RATE

Soil application

IN-FURROW OR BANDED AT PLANTING OR SIDE-DRESS: Apply 1.5-3 fl. oz. per acre. A follow-up foliar application can be made during early vegetative growth stages.

BROADCAST: Apply 3-5 fl. oz. per acre. Ensure sufficient rain or irrigation to place the product in the root zone.

DRIP OR MICRO IRRIGATION: Apply 3-5 fl. oz. per acre. For soil application to irrigated crops, follow the same application schedule as described with foliar applications.

Foliar application

VINES: Apply 3 applications at 3-6 fl. oz. per acre respectively during pre-bloom vegetative stage, during flowering to cap fall, and again 21-28 days later (pre-veraison). Start at the low rate and increase the rate with each application as the foliage expands. For red table grape varieties, a fourth application can be made at the start of veraison at 5-6 fl. oz. per acre. A post-harvest application can be considered at 6 fl. oz. per acre within 2 weeks after harvest. Use a minimum water volume of 40 gallons per acre.

DECIDUOUS TREES: Apply 3 applications at 3-5 fl. oz. per acre per application, starting at 50-100% bloom on pome fruit, or at 75-100% petal drop on stone fruit and nuts. Repeat 2 more applications with 21–28-day intervals. Start at the low rate and increase the rate with each application as the foliage expands. Do not apply closer than 1 month from harvest. A post-harvest application can be considered at 5 fl. oz. per acre within 2 weeks after harvest. Use a minimum water volume of 40 gallons per acre.

Foliar application, continued

CITRUS, AVOCADOS, AND OTHER SUBTROPICAL TREES: Apply 3-4 applications at 5.5 fl. oz. per acre, each starting at first growth flush in spring, 75-90% petal drop, and repeat with 28-day intervals up to 4 applications. Use a minimum water volume of 75 callons per acre.

BERRIES AND SMALL FRUITS: Apply 2-4 fl. oz. per acre per application, 3-4 applications, 21-28-day intervals, during vegetative growth, early bloom, and after fruit set. Use a minimum water volume of 25 gallons per acre.

NON-BEARING TREES: Apply 6 fl. oz. per 100 gallons (0.05% v/v), 3-4 applications, 21-28-day intervals, starting within 2 weeks of spring flush.

VEGETABLES: Apply 1.5-4 fl. oz. per acre per application starting at the low rate and increasing the rate as the crop canopy grows, 3-5 applications, 21-28-day intervals, starting 14 days after germination or transplant, continuing through flowering and fruit development or tuber or bulb growth. Use a minimum water volume of 20 gallons per acre

VEGETABLE SEEDLINGS: Spray or drench vegetable seedlings twice with 14-day intervals prior to transplant with a solution of 7 fl. oz. per 100 gal (2 ml per gallon).

ROW CROPS: Apply 1.5-2 fl. oz. per acre during early vegetative growth. A second application at 2 fl. oz. per acre can be applied during flag leaf on cereals or at R1-R3 stages on flowering crops. Use a minimum water volume of 15 gallons per acre.

STORAGE AND DISPOSAL

STORAGE: Keep away from open flames, hot surfaces and sources of ignition. Store in a cool, dry, well-ventilated place out of direct sunlight. Store in original tightly closed container away from children, animals, foods, feeds and seeds. PRODUCT AND CONTAINER DISPOSAL: Do not allow this material to drain into sewers/water supplies. Empty containers or liners may retain some product residues. Triple rinse (or equivalent) during mixing and loading. Dispose of contents/container in accordance with local/regional/national/international regulations.

NOTICE OF WARRANTY

CONDITIONS OF SALE-LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES: To be read prior to using this product. The directions should be followed carefully together with good application practice. Failure to do so may cause injury to crops, animals, man or the environment. Extraordinary or unusual weather conditions may affect product performance and/or crop injury which are beyond the control of Oro Agri, Inc. (the "Company") or seller. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions For Use. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law. The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Company's election, ONE of the following: 1. Refund of the purchase price paid by buyer or user for product bought. 2. Replacement of the product used. To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The company/seller offer this product and the buyer/user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

For more information contact info@oroagri.rovensa.com.

GALACTEK, WETCIT, and VINTRE are proprietary trademarks of Oro Agri, Inc. v20240528