\*(1) Acid products are for use as a supplemental fertilizer treatment. \*(2) Foliar application, the phosphite ions are taken up directly by the plant foliage and may undergo a degree of conversion to phosphate ions, or will be used directly by plants as phosphita ions.

"(3) Soil application to annual crops, a lesser response from the initial crop, with a corresponding superior response from succeeding crops, may be observed. In addition, placement close to seeds or root zones may be injurious to crops. The effect may be aggravated by a soil pH below 6.5.

Item# 5492-CA

# Sysstem-LeafMax<sup>™</sup> 0-20\*-0



### Warning.

Causes skin and eye irritation, harmful if swallowed, may be harmful to aquatic life.

Avoid breathing mist. Wash thoroughly after handling. Use only outdoors or in self-vertilated area. Avoid release to aquatic environment. Wear protective gloves/clothing/eyewear/face. If on skin: Wash with plenty of soap and water. If inhaled: Remove person to fresh air and keep comfortable for breathing. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do, continue inising. Call a poison center/doctor fly out feel unwell. If skin or eye irritation occurs: Set medical advice/attention. Take off contaminated clothing and wash before reuse. Dispose of contents/ container in accordance with local/regional/indional regulations.

Agro-K Corporation 8030 Main Street, N.E. • Minneapolis, MN • 55432-1844 U.S.A. (800) 328-2418 • (763) 780-4116 • www.agro-k.com

# Science-Driven Nutrition<sup>SM</sup>

# Directions for Commercial Use:

Foliar: Apply 2-8 pints per acre with sufficient water for thorough coverage. Product can be applied at 7 to 14 day intervals for a maximum of 5 times during the growing season.

Soit: Apply 1 to 4 quarts per acre. Product can be applied at 7 to 14 day intervals for a maximum of 5 times during the growing season. Can be band sprayed or metered into irrigation system. Product can also be applied as a soil drench on trees.

Foliar fertilization is intended to supplement standard ground fertility programs and will not by itself provide all nutrients normally required by agricultural crops.

### APPLICATION PRECAUTIONS:

### Tank Mixes

Sysstem-Leaf Max is compatible with many fertilizers and pesticides but a compatibility check is always recommended. Fill the tank to 26 year, but water and begin agitation and add tank-mix partner products (surfactants, pesticides, etc.) to the tank based on manufacturer labeled instructions. Mixing order will depend on the tank-mix partners. If the tank-mix partners are esensitive to high pH, add Sysstem-Leaf Max first to lower the pH of the spray solution. If the tank-mix partners are sensitive to low pH solutions, add Sysstem-Leaf Max is and spray immediately after mixing. For adjuvants refer the tank-mix partner labels for the proper mixing order and adjuvant selection. Do not mix Sysstem-Leaf Max with fixed copper-based fungicides. Do not apply Sysstem-Leaf Max in combination with citrus oils at more than 1 pt/ac or 0.5%, whichever is less or in combination with petroleum oils at more than 2 dtx/ac or 1.0% whichever is less or in combination with petroleum oils at more than 2 dtx/ac or 1.0% whichever is less.

**WARNING:** The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

### Guaranteed Analysis:

Total Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	20.0%
Magnesium (Mg)	0.50%
Sulfur (S)	2.0%
Cobalt (Co)	0.005%
Copper (Cu)	0.50%
Iron (Fe)	0.75%
Manganese (Mn)	1.25%
Molybdenum (Mo)	0.15%
Zinc (Zn)	3.0%

### Derived from

Phosphorous Acid, Cobalt Carbonate, Copper Sulfate, Ferrous Sulfate, Magnesium Carbonate, Manganese Carbonate, Sodium Molybdate. Zinc Carbonate

Product density is 10.84 lbs/gallon at 68 degrees Fahrenheit.

Net Wgt 27.10 lbs / 2.5 gal Net Wgt 12.30kg / 9.46L



# Spray Applications

Do not apply Sysstem-Leaf Max with any type of hand, pump or back-pack sprayer. Avoid applying to wet leaf and fruit surfaces. Apply to dry plant surfaces to the point of wet with no run-off. Exercise caution spraying when temperature is above 92°F, do not apply in extreme temperature. For additional information regarding Agro-K products, tank-mixes and spray conditions contact your local Agro-K representative. <u>User assumes full responsibility to ensure product, equipment and plant compatibilities.</u>

**WARRANTY** - 1.-The manufacturer guarantees and warrants that the content and the total net weight are as stated within lawful limits.

2.-Liability of Agro-K Corporation under this warranty or otherwise shall be limited to refund of the purchase price and such refund is expressly agreed by the buyer to be the exclusive remedy.

Agro-K Corporation makes no other expressed or implied guarantee, warranty or representation, including warranties of merchantability and fitness for a purpose. Agro-K Corporation shall not be liable for direct consequential or incidental damages. No modifications of this warranty and the disclaimers herein are authorized or valid unless expressed in writing and signed by Agro-K Corporation.

### SHAKE WELL BEFORE USE WARNING: KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

V1